

2023-2024 学年第一学期期中考试卷

Khartoum: ٢٠٢٢

گزٹوم فن

No/Min/O.H/:

النمرة: وص / موص /

Sudan Federal Ministry of Health

Primary Health Care Directorate

The National Model Curriculum Project

and Soudan. This is the first time that AI has been used to predict the risk of flooding in the Blue Nile basin.

The results of the study have been presented at the 2019 African Floods Conference in Addis Ababa, Ethiopia, under the title "A hydrological model for flood prediction in the Blue Nile basin using AI".

The study was conducted by a team of researchers from the University of Khartoum, the National Water Research Institute (NWI) in Canada, and the International Centre for Environmental and Nuclear Sciences (ICENS) in Saudi Arabia.

The study found that AI can be used to predict the risk of flooding in the Blue Nile basin, which is one of the most vulnerable regions in Africa to flooding.

The study also found that AI can be used to predict the risk of flooding in the Blue Nile basin, which is one of the most vulnerable regions in Africa to flooding.

The study also found that AI can be used to predict the risk of flooding in the Blue Nile basin, which is one of the most vulnerable regions in Africa to flooding.

The study also found that AI can be used to predict the risk of flooding in the Blue Nile basin, which is one of the most vulnerable regions in Africa to flooding.

The study also found that AI can be used to predict the risk of flooding in the Blue Nile basin, which is one of the most vulnerable regions in Africa to flooding.

The study also found that AI can be used to predict the risk of flooding in the Blue Nile basin, which is one of the most vulnerable regions in Africa to flooding.

The study also found that AI can be used to predict the risk of flooding in the Blue Nile basin, which is one of the most vulnerable regions in Africa to flooding.

The study also found that AI can be used to predict the risk of flooding in the Blue Nile basin, which is one of the most vulnerable regions in Africa to flooding.

The study also found that AI can be used to predict the risk of flooding in the Blue Nile basin, which is one of the most vulnerable regions in Africa to flooding.

The study also found that AI can be used to predict the risk of flooding in the Blue Nile basin, which is one of the most vulnerable regions in Africa to flooding.

The study also found that AI can be used to predict the risk of flooding in the Blue Nile basin, which is one of the most vulnerable regions in Africa to flooding.

The study also found that AI can be used to predict the risk of flooding in the Blue Nile basin, which is one of the most vulnerable regions in Africa to flooding.

The study also found that AI can be used to predict the risk of flooding in the Blue Nile basin, which is one of the most vulnerable regions in Africa to flooding.

The study also found that AI can be used to predict the risk of flooding in the Blue Nile basin, which is one of the most vulnerable regions in Africa to flooding.

The study also found that AI can be used to predict the risk of flooding in the Blue Nile basin, which is one of the most vulnerable regions in Africa to flooding.

In 2005, the Government of Sudan, represented by the Federal Ministry of Health

The distribution of the drug started at the beginning of 2005. The first batch of 100,000 tablets was distributed to the State of Kordofan, followed by 100,000 tablets to the State of North Darfur. These distributions were followed by others to other states such as South Darfur, Central Darfur, West Darfur, and the State of Al-Jazira.

After the distribution of the drug, the National Center for Disease Control (Ministry of Health) organized training sessions for health workers in all the states where the drug was distributed.

Training sessions were organized for health workers in all the states where the drug was distributed. These sessions were organized by the National Center for Disease Control (Ministry of Health).

Training sessions were organized for health workers in all the states where the drug was distributed. These sessions were organized by the National Center for Disease Control (Ministry of Health).

Training sessions were organized for health workers in all the states where the drug was distributed. These sessions were organized by the National Center for Disease Control (Ministry of Health).

Training sessions were organized for health workers in all the states where the drug was distributed. These sessions were organized by the National Center for Disease Control (Ministry of Health).

Training sessions were organized for health workers in all the states where the drug was distributed. These sessions were organized by the National Center for Disease Control (Ministry of Health).

In 1995 the prevalence was 3%

In 1995 the prevalence was 3%. In 2005, the prevalence was 1%. This indicates that the prevalence has decreased significantly over the past decade.


c) In 2007 the prevalence was 0.45%.

d) In 2011 the prevalence was 0.00%.

The allowed prevalence is 0.5% or less.

100% 90% 80% 70% 60% 50% 40% 30% 20% 10% 0%

0.5% 0.45% 0.40% 0.35% 0.30% 0.25% 0.20% 0.15% 0.10% 0.05% 0.00%



4. The Lions Club International Foundation's Contribution to the Cancer Society

